

Text for INOPIL Insert

Boni

Calcium 500 mg

Boni is a preparation of Calcium Carbonate. Calcium is used in the treatment of deficiency states and as a dietary supplement when intake of calcium may be inadequate. Calcium Carbonate reacts with gastric acid to produce Calcium chloride and water. From this Calcium chloride, elemental calcium is absorbed from gastrointestinal tract.

COMPOSITION

Each tablet contains Calcium Carbonate BP 1250 mg equivalent to elemented Calcium 500 mg.

INDICATION

Calcium Carbonate is used for the treatment or prevention of calcium depletion in patients in whom dietary measures are inadequate. Conditions that may be associated with calcium deficiency include hypoparathyroidism, achlorhydria, chronic diarrhoea, vitamin D deficiency, steatorrhea, spree, pregnancy and lactation, menopause, pancreatitis, renal failure, alkalosis and hyperphosphatemia. Calcium Carbonate is being used increasingly to treat hyperphosphatemia in chronic renal failure as well as those on continuous ambulatory peritoneal dialysis (CAPD) and haemodialysis. Calcium Carbonate containing preparation can provide short term relief of dyspeptic system but are no longer recommended for long-term treatment of peptic ulceration. It is also indicated for prevention and treatment of osteoporosis, osteomalacia, disorders of osteogenesis & tooth formation and latent teary.

DOSAGE & ADMINISTRATION

One tablet daily or as directed by the physician. Higher doses should not be taken unless recommended by the physician.

SIDE EFFECT

In rare cases, gastrointestinal disturbances e.g. diarrhoea, constipation & flatulence, hypercaemia, bradycardia and arrhythmia.

CONTRAINDICATION

Boni is contraindicated to hypercalcaemia, nephrolithiasis, hyperparathyroidism and Zollinger-Ellison Syndrome.

PRECAUTION

Calcium salts should be used cautiously in patients with sarcoidosis, renal or cardiac diseases and in patients receiving cardiac glycosides. When hypercalcaemia occurs, discontinuation of the drug is usually sufficient to return serum calcium concentration to normal.

PREGNANCY AND LACTATION

Calcium containing drugs have been widely used in pregnancy by way of oral calcium supplementation or antacid therapy. It can be used in lactating women too.

DRUG INTERACTION

Calcium Carbonate may enhance the cardiac effects of digoxin and other cardiac glycosides. Calcium Carbonate may interfere with the absorption of concomitantly administered tetracycline.

PACKAGING

Box containing 5x10 tablets in blister pack.

PHARMACEUTICAL PRECAUTION

Keep in a dry place away from light and heat.

WARNING

Keep out of the reach of children.

Manufactured by



Delta Pharma Limited
Kishoreganj, Bangladesh